

## Synaptogenesis in the Rat Cerebellum: Effects of Early Hypo- and Hyperthyroidism

**Abstract.** *The number of synapses in the molecular layer of the rat cerebellum is reduced by early hypo- and hyperthyroidism within 30 days. Hypothyroidism retards synaptogenesis after 10 days, while hyperthyroidism accelerates synaptogenesis initially, but by 21 days the number of synapses is reduced. The sensitivity of developing synapses to thyroid hormone may permit analysis of the events triggering synaptogenesis.*

Neonatal hypo- and hyperthyroidism produce various deficits in postnatal neural development, including a decrease in cerebral and cerebellar weights (1, 2), and changes in the number, size, and packing density of cells (1-3). Changes in the amount and composition of the neuropil have been demonstrated in the sensorimotor cortex (2, 4), the visual cortex (5), and the cerebellum (6). Changes have also been found in development of metabolic compartmentation, which is thought to reflect maturation of dendritic processes and nerve terminals (7). Together these results indicate a retardation of neuropil development in hypothyroidism, and an acceleration in hyperthyroidism. Also, there is behavioral and electrophysiological evidence for neurological changes in these conditions in the form of retarded or accelerated maturation of innate behavioral patterns (8, 9), and abnormalities in the electrical patterns of the brain (8, 10).

We examined the effects of early hypo- and hyperthyroidism on synaptogenesis in the cerebellar molecular layer with quantitative light and electron microscope methods. Our results provide evidence that both hypo- and hyperthyroidism cause a reduction in the total number of synapses formed in the cerebellar molecular layer, but by different processes.

Groups of animals were injected




Fig. 1. Density of synaptic profiles in cerebellar molecular layer.  $\Delta$ , Hypert thyroid;  $\circ$ , hypo thyroid;  $\bullet$ , control. Statistical significance: control and hypo thyroid, at 21 and 30 days,  $P < .01$ ; at 55 days,  $P < .05$ ; control and hypert thyroid, at 10, 15, and 21 days,  $P < .01$ .

from birth with either physiological saline (controls), propylthiouracil (producing a hypothyroidism), or L-thyroxine (producing hyperthyroidism) (11). Each litter was composed of ten rats (mostly males) originating from at least two different litters born on the same day. Animals were killed at 5, 10, 15, 21, 24, or 30 days of age. For light microscopy, the brains were fixed in Bouin's fluid followed by 10 percent neutral formalin, and were embedded in Paraplast. The brains were sectioned sagittally at 6  $\mu\text{m}$  and parasagittal sections were stained with hematoxylin and eosin. The area of the molecular layer was determined with an Ott compensating polar planimeter applied to cerebellar tracings of these sections (magnified 65 times with a modified Leitz projection apparatus). For electron microscopy, the brains were fixed by perfusion with 6 percent glutaraldehyde buffered with phosphate (pH 7.2). Cerebellums were removed and were further fixed in cold 6 percent glutaraldehyde for at least 1 hour. This was followed by slicing at 235  $\mu\text{m}$ , dehydration in ethanol, and staining with ethanolic phosphotungstic acid (12), after which tissue slices were embedded in Araldite-Epon. Samples for synaptic density counts were randomly selected from outer, middle, and inner zones of the molecular layers from coded animals. Pictures were taken at a magnification of  $\times 6027$  with a Philips 300 electron microscope. Synapse counts were made directly from coded negatives and expressed as the average number of synaptic profiles per 96  $\mu\text{m}^2$  of molecular layer. To obtain estimates of the total number of synapses in the sagittal area of the molecular layer, we applied the formula: total synaptic profiles equals synaptic profiles per 96  $\mu\text{m}^2$  of the molecular layer times the area of molecular layer in square micrometers. Analysis of variance and Duncan's multiple range test (13) were used as tests of significance ( $P < .01$ ).

Early hypothyroidism caused re-

tardation of synaptogenesis as shown by a reduced rate of increase in the density of synaptic profiles; this was significant at 21, 30, and 55 days (Fig. 1). A similar pattern was seen in the development of the molecular layer (Fig. 2). By 30 days the area of the molecular layer was still significantly reduced, although not as much as in the hyperthyroid group. The increase in the calculated total number of synapses was retarded from day 10 on, but by day 30, the reduction was not as great as in the hyperthyroid group.

Early hyperthyroidism caused a transient increase in density of synaptic profiles until 24 days of age followed by a decline to control values by 30 days (Fig. 1). This treatment also caused a pronounced decrease in area of the molecular layer after 15 days (Fig. 2). The calculated total number of synapses in the hyperthyroid animals was higher than in the controls until 21 days, followed by a significant reduction at 30 days as compared with controls.

Our results show that hypo- and hyperthyroidism lead to a pronounced reduction in the synaptic content of the cerebellar molecular layer. However, this effect at day 30 is probably the result of two different processes. In hyperthyroidism, the cells of the external granular layer (postnatal germinal matrix of the cerebellum) have been shown to cease proliferation early (14) producing fewer stem cells from which granule, basket, and stellate cells are formed. This premature termination is associated with early initiation of cell differentiation (14) leading to an initial acceleration of synaptogenesis but to an ultimate reduction in total number of synapses. In hypothyroidism,




Fig. 2. Area of cerebellar molecular layer.  $\Delta$ , Hypert thyroid;  $\circ$ , hypo thyroid;  $\bullet$ , control. Statistical significance: control and hypo thyroid, 15 to 30 days,  $P < .01$ ; control and hypert thyroid, 21 to 30 days,  $P < .01$ .




Fig. 3. Estimated total synaptic profiles in cerebellar molecular layer.  $\Delta$ , Hyperthyroid;  $\circ$ , hypothyroid;  $\bullet$ , control.

however, the main effect seems to be a general retardation of the differentiation of cerebellar neurons (6, 14, 15). The evidence of retarded synaptogenesis throughout development, shown in this study, supports this hypothesis.

In conclusion, it appears from these and other results (14) that both the acceleration of cell differentiation in the cerebellar cortex (produced by hyperthyroidism) and its retardation (produced by hypothyroidism) lead to reductions in the synaptic content of the neuropil. It remains to be determined whether or not acceleration or retardation produced by other means, and in other parts of the brain, also lead to synaptic deficits. These results not only suggest the possible role of thyroid hormone as a trigger in the process of synaptogenesis, but also indicate the usefulness of early thyroid treatments as a tool for studying synaptogenesis in the developing nervous system.

JEAN L. NICHOLSON

JOSEPH ALTMAN

Laboratory of Developmental  
Neurobiology, Department of  
Biological Sciences, Purdue  
University, Lafayette, Indiana 47906

#### References and Notes

- R. Balázs, S. Kovács, P. Teichgraber, W. A. Cocks, J. T. Eayrs, *J. Neurochem.* **15**, 1335 (1968); R. Balázs, S. Kovács, W. A. Cocks, A. L. Johnson, J. T. Eayrs, *Brain Res.* **25**, 555 (1971).
- J. T. Eayrs and S. H. Taylor, *J. Anat.* **85**, 350 (1951).
- S. E. Geel and P. S. Timiras, *Endocrinology* **80**, 1069 (1967); J. M. Pasquini, B. Kaplin, C. A. García Argiz, C. J. Gomez, *Brain Res.* **6**, 621 (1967); L. Krawiec, C. A. García Argiz, C. J. Gomez, J. M. Pasquini, *ibid.* **15**, 209 (1969).
- J. T. Eayrs and G. Horn, *Anat. Rec.* **121**, 53 (1955); J. T. Eayrs, *Growth* **25**, 175 (1961).
- B. G. Cragg, *Brain Res.* **18**, 297 (1970).
- J. P. Geloso, P. Herson, J. Legrand, C. Legrand, A. Jost, *Gen. Comp. Endocrinol.* **10**, 191 (1968); G. Lefranc, Y. George, J. Tusques, *C.R. Soc. Biol.* **162**, 219 (1968); J. Legrand, *C.R.H. Acad. Sci.* **261**, 544 (1965); *Arch. Anat. Microsc. Morphol. Exp.* **56**, 206 (1967); in *Regional Development of the Brain in Early Life*, A. Minkowski, Ed. (Blackwell, Oxford, 1967); J. Tusque, G. Lefranc, Y. George, *C.R. Soc. Biol.* **161**, 2256 (1967).
- J. A. Cocks, R. Balázs, A. L. Johnson, J. T. Eayrs, *J. Neurochem.* **17**, 1275 (1970); A. J. Patel and R. Balázs, *ibid.* p. 955.
- S. Schapiro and R. J. Norman, *Science* **155**, 1279 (1967).
- J. T. Eayrs, *Anim. Behav.* **12**, 195 (1964); M. Hamburgh and E. Vicari, *Anat. Rec.* **127**, 302 (1957).
- D. M. Woodbury, R. E. Hurley, W. G. Lewis, M. W. Arthur, W. W. Copeland, J. F. Kirschvink, L. S. Goodman, *J. Pharmacol. Exp. Ther.* **106**, 331 (1952); P. B. Bradley, J. T. Eayrs, K. Schmalbach, *Electroencephalogr. Clin. Neurophysiol.* **12**, 467 (1960); M. Salas and S. Schapiro, *Physiol. Behav.* **5**, 17 (1970).
- Physiological saline: 0.05 ml on days 0 to 7; 0.1 ml on days 8 to 30. Propylthiouracil (PTU): 0.05 ml of 0.2 percent PTU (in 1 percent carboxymethyl-cellulose) on days 0 to 10; 0.1 ml on days 11 to 20; 0.1 ml of 0.4 percent PTU on days 21 to 30. L-Thyroxine: 1  $\mu$ g (in physiological saline) on days 0 to 7; 2  $\mu$ g on days 8 to 14; 3  $\mu$ g on days 15 to 21; 5  $\mu$ g on days 22 to 30 (16). The extent of hypothyroidism caused by PTU was determined by histologically monitoring the thyroid for lack of colloid and hypoplastic follicular epithelium; PTU caused complete blockage as early as day 5.
- F. E. Bloom and G. K. Aghajanian, *J. Ultrastruct. Res.* **22**, 361 (1968).
- D. B. Duncan, *Biometrika* **11**, 1 (1955).
- J. Nicholson and J. Altman, *Brain Res.*, in press.
- M. Hamburgh, *Gen. Comp. Endocrinol.* **10**, 198 (1968); ———, L. A. Mendoza, J. F. Burkhardt, F. Weil, in *Hormones in Development*, M. Hamburgh and E. J. W. Barrington, Eds. (Appleton-Century-Crofts, New York, 1971), pp. 403-415.
- M. Hamburgh, E. Lynn, E. P. Weiss, *Anat. Rec.* **150**, 147 (1964).
- Supported by NIH and AEC.

30 September 1971; revised 18 January 1972